



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: David M. Lubman *et al.*

Serial No.: 09/778,548

Filed: 02/07/01

Entitled: Protein Separation And Display

Group No.: 1631

Examiner: Marschel

INFORMATION DISCLOSURE
STATEMENT TRANSMITTAL

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

CERTIFICATE OF MAILING UNDER 37 C.F.R. § 1.8(a)(1)(i)(A)

I hereby certify that this correspondence (along with any referred to as being attached or enclosed) is, on the date shown below, being deposited with the U.S. Postal Service with sufficient postage as first class mail in an envelope addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Dated: October 10, 2003

By: Mary Ellen Waite

Mary Ellen Waite

Sir or Madam:

Enclosed please find an Information Disclosure Statement and Form PTO-1449, including copies of the references contained thereon, for filing in the U.S. Patent and Trademark Office.

A check for \$180.00 is also enclosed pursuant to 37 C.F.R. § 1.17(p) for filing this Information Disclosure Statement after three months as set forth in 37 C.F.R. § 1.97(c).

The Commissioner is hereby authorized to charge any additional fee or credit overpayment to our Deposit Account No. 08-1290. **An originally executed duplicate of this transmittal is enclosed for this purpose.**

Dated: October 10, 2003

Tanya A. Arenson

Tanya A. Arenson
Registration No. 47,391

MEDLEN & CARROLL, LLP
101 Howard Street, Suite 350
San Francisco, California 94105
617/252-3353

RECEIVED
OCT 17 2003
TECH CENTER 1600/2900



PATENT
Attorney Docket No. **UM-06102**

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: David M. Lubman *et al.*
Serial No.: 09/778,548
Filed: 02/07/01
Entitled: **Protein Separation And Display**

Group No.: 1631
Examiner: A.H. Marschel

INFORMATION DISCLOSURE STATEMENT

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

CERTIFICATE OF MAILING UNDER 37 C.F.R. § 1.8(a)(1)(i)(A)	
I hereby certify that this correspondence (along with any referred to as being attached or enclosed) is, on the date shown below, being deposited with the U.S. Postal Service with sufficient postage as first class mail in an envelope addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.	
Dated: <u>October 10, 2003</u>	By: <u>Mary Ellen Waite</u> Mary Ellen Waite

Sir or Madam:

The citations listed below, copies attached, may be material to the examination of the above-identified application, and are therefore submitted in compliance with the duty of disclosure defined in 37 C.F.R. §§ 1.56 and 1.97. The Examiner is requested to make these citations of official record in this application.

10/16/2003 MROCHA1 00000012 09778548

01 FC:1886

188.00 CP

The following U.S. patent applications are closely related to this specification (not provided):

- 20020039747 (09/778496), Lubman, et al., filed 02/07/01, Mapping of Differential Display of Proteins;
- 20030064527 (10/133896), Lubman, *et al.*, filed 04/26/02, Proteomic Differential Display;
- 20030058239 (10/133711), Wall, *et al.*, filed 04/26/02, Three-Dimensional Protein Mapping;
- 09/968930, Lubman, *et al.*, filed 10/1/01, Protein Mapping; and
- 10/448,564 Lubman, *et al.*, 5/30/03, Automated Protein Analysis System.

The following PCT patents and applications are closely related to this application:

- WO0158925, 08/16/01, CHONG *et al.*, Protein Separation and Display;
 - WO02088701, 11/07/02, PARUS *et al.*, Methods of Multi-Phase Protein Analysis;
 - WO0158926, 03/2/03, CHONG *et al.*, Protein Mapping;
 - WO0159460, 03/2/03, CHONG *et al.*, Mapping of Differential Display of Proteins;
- and
- PCT/US03/16899, 5/30/03, PARUS *et al.*, 5/30/03, Automated Protein Analysis System.

The following printed publications are referred to in the body of the specification:

- U.S. 6002127 Vestal, 12/14/99, Time-of-Flight Mass Spectrometry Analysis of Biomolecules;
- Ayala *et al.*, Use of Rotofor preparative isoelectrofocusing cell in protein purification procedure, Appl Biochem Biotechnol, 69(1):11-16 (1998);

- Bini *et al.*, Protein expression profiles in human breast ductal carcinoma and histologically normal tissue, *Electrophoresis*, 18(15):2832-2841 (1997);
- Chen *et al.*, Identification of proteins from two-dimensional gel electrophoresis of human erythroleukemia cells using capillary high performance liquid chromatography/electrospray-ion trap-reflectron time-of-flight mass spectrometry with two-dimensional topographic map analysis of in-gel tryptic digest products, *Rapid Commun Mass Spectrom*, 13(19):1907-1916 (1999);
- Chen *et al.*, Rapid identification and screening of proteins from whole cell lysates of human erythroleukemia cells in the liquid phase, using non-porous reversed phase high-performance liquid chromatography separations of proteins followed by matrix-assisted [correction of multi-assisted] laser desorption/ionization mass spectrometry analysis and sequence database searching, *Rapid Commun Mass Spectrom*, 12(24):1994-2003 (1998);
- Chong *et al.*, Rapid screening of protein profiles of human breast cancer cell lines using non-porous reversed-phase high performance liquid chromatography separation with matrix-assisted laser desorption/ionization time-of-flight mass spectral analysis, *Rapid Commun Mass Spectrom*, 13(18):1808-1812 (1999);
- Dai *et al.*, Two-layer sample preparation: a method for MALDI-MS analysis of complex peptide and protein mixtures, *Anal Chem*, 71(5):1087-1091 (1999);
- Figey and Aebersold, Microfabricated modules for sample handling, sample concentration and flow mixing: application to protein analysis by tandem mass spectrometry, *J Biomech Eng*, 121(1):7-12 (1999);
- Fuqua *et al.*, Induction of the estrogen-regulated "24K" protein by heat shock, *Cancer Res*, 49(15):4126-4129 (1989);
- Griffin *et al.*, Direct database searching with MALDI-PSD spectra of peptides, *Rapid Commun Mass Spectrom*, 9(15):1546-1551 (1995);
- Hanash, *Advances in Electrophoresis*, 1-44, (1998);

- Herbert, Advances in protein solubilisation for two-dimensional electrophoresis, *Electrophoresis*, 20(4-5):660-663 (1999);
- Immler *et al.*, Identification of phosphorylated proteins from thrombin-activated human platelets isolated by two-dimensional gel electrophoresis by electrospray ionization-tandem mass spectrometry (ESI-MS/MS) and liquid chromatography-electrospray ionization-mass spectrometry (LC-ESI-MS), *Electrophoresis*, 19(6):1015-1023 (1998);
- Liang *et al.*, Determination of bacterial protein profiles by matrix-assisted laser desorption/ionization mass spectrometry with high-performance liquid chromatography, *Rapid Commun Mass Spectrom*, 10(10):1219-1226 (1996);
- Matsui *et al.*, Immobilized pH gradient two-dimensional gel electrophoresis and mass spectrometric identification of cytokine-regulated proteins in ME-180 cervical carcinoma cells, *Electrophoresis*, 18(3-4):409-417 (1997);
- Mohammad *et al.*, Induced expression of alpha-enolase in differentiated diffuse large cell lymphoma, *Enzyme Protein*, 48(1):37-44 (1994);
- Neubauer and Mann, Mapping of phosphorylation sites of gel-isolated proteins by nanoelectrospray tandem mass spectrometry: potentials and limitations, *Anal Chem*, 71(1):235-242 (1999);
- Nilsson *et al.*, Isolation and characterization of proteins from human lymphocyte nuclei using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry and post-source decay analysis, *Rapid Commun Mass Spectrom*, 11(6):610-612 (1997);
- Opiteck *et al.*, Comprehensive two-dimensional high-performance liquid chromatography for the isolation of overexpressed proteins and proteome mapping, *Anal Biochem*, 258(2):349-361 (1998);
- Patterson and Aebersold, Mass spectrometric approaches for the identification of gel-separated proteins, *Electrophoresis*, 16(10):1791-1814 (1995);
- Pinto *et al.*, An enhanced microfluidic chip coupled to an electrospray Qstar mass

spectrometer for protein identification, *Electrophoresis*, 21(1):181-190 (2000);

- Rasmussen *et al.*, Two-dimensional gel database of human breast carcinoma cell expressed proteins: an update, *Electrophoresis*, 19(5):818-825 (1998);
- Redner *et al.*, The t(5;17) variant of acute promyelocytic leukemia expresses a nucleophosmin-retinoic acid receptor fusion, *Blood*, 87(3):882-886 (1996);
- Reid *et al.*, Capillary column chromatography improves sample preparation for mass spectrometric analysis: complete characterization of human alpha-enolase from two-dimensional gels following in situ proteolytic digestion, *Electrophoresis*, 19(6):946-955 (1998);
- Righetti, Isoelectric focusing : theory, methodology, and applications, Elsevier Biomedical Press; Elsevier-North Holland, 1st, (1983);
- Rosenfeld *et al.*, In-gel digestion of proteins for internal sequence analysis after one- or two-dimensional gel electrophoresis, *Anal Biochem*, 203(1):173-179 (1992);
- Sanchez *et al.*, Inside SWISS-2DPAGE database, *Electrophoresis*, 16(7):1131-1151 (1995);
- Sirover, New insights into an old protein: the functional diversity of mammalian glyceraldehyde-3-phosphate dehydrogenase, *Biochim Biophys Acta (BBA) - Protein Structure and Molecular Enzymology*, 1432(2):159-184 (1999);
- Steller, Mechanisms and genes of cellular suicide, *Science*, 267(5203):1445-1449 (1995);
- ten Hoeve *et al.*, Isolation and chromosomal localization of CRKL, a human crk-like gene, *Oncogene*, 8(9):2469-2474 (1993);
- Wall *et al.*, Isoelectric focusing nonporous RP HPLC: a two-dimensional liquid-phase separation method for mapping of cellular proteins with identification using MALDI-TOF mass spectrometry, *Anal Chem*, 72(6):1099-1111 (2000);
- Wall *et al.*, Rapid profiling of induced proteins in bacteria using MALDI-TOF mass

spectrometric detection of nonporous RP HPLC-separated whole cell lysates, *Anal Chem*, 71(17):3894-3900 (1999);

- Welsh *et al.*, Variation in expression of hsp27 messenger ribonucleic acid during the cycle of the seminiferous epithelium and co-localization of hsp27 and microfilaments in Sertoli cells of the rat, *Biol Reprod*, 55(1):141-151 (1996);
- Yates, Mass spectrometry and the age of the proteome, *J Mass Spectrom*, 33(1):1-19 (1998); and
- Zugaro *et al.*, Characterization of rat brain stathmin isoforms by two-dimensional gel electrophoresis-matrix assisted laser desorption/ionization and electrospray ionization-ion trap mass spectrometry, *Electrophoresis*, 19(5):867-876 (1998).

The following additional references were cited in International Search Reports for the corresponding PCT application WO0158925:

- Chong *et al.*, Rapid profiling of E. coli proteins up to 500 kDa from whole cell lysates using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, *Rapid Commun Mass Spectrom*, 11(17):1900-1908 (1997);
- Opiteck *et al.*, Two-dimensional SEC/RPLC coupled to mass spectrometry for the analysis of peptides, *Anal Chem*, 69(13):2283-2291 (1997); and
- Regnier and Huang, Future potential of targeted component analysis by multidimensional liquid chromatography-mass spectrometry, *J Chromatogr A*, 750(1-2):3-10 (1996);
- Andrews *et al.*, Analysis of DNA adducts using high-performance separation techniques coupled to electrospray ionization mass spectrometry, *J Chromatogr A*, 856(1-2):515-526 (1999);
- Ball, Purification of synthetic peptides with the aid of reversible chromatographic probes, *J Chromatogr A*, 686(73-83 (1994);

- Binz *et al.*, A molecular scanner to automate proteomic research and to display proteome images, *Anal Chem*, 71(21):4981-4988 (1999);
- Davidsson and Nilsson, Peptide mapping of proteins in cerebrospinal fluid utilizing a rapid preparative two-dimensional electrophoretic procedure and matrix-assisted laser desorption/ionization mass spectrometry, *Biochim Biophys Acta*, 1473(2-3):391-399 (1999);
- Lee, Protein separation using non-porous sorbents, *J Chromatogr B Biomed Sci Appl*, 699(1-2):29-45 (1997);
- Medzihradszky *et al.*, Protein sequence and structural studies employing matrix-assisted laser desorption ionization-high energy collision-induced dissociation, *International Journal of Mass Spectrometry and Ion Processes*, 160(1-3):357-369 (1997);
- Nimura *et al.*, Fast protein separation by reversed-phase high-performance liquid chromatography on octadecylsilyl-bonded non-porous silica gel. Effect of particle size of column packing on column efficiency., *J Chromatogr*, 585(207-211 (1991);
- Richmond, High-throughput flow injection analysis-mass spectrometry with networked delivery of colour rendered results: the characterisation of liquid chromatography fractions, *J Chromatogr A*, 835(29-39 (1999);
- Wilkins *et al.*, From proteins to proteomes: large scale protein identification by two-dimensional electrophoresis and amino acid analysis, *Biotechnology (N Y)*, 14(1):61-65 (1996);
- U.S. 5498545, 3/12/96, Vestal, Mass Spectrometer System and Method for Matrix-Assisted Laser Desorption Measurements;
- U.S. 5670054, 9/23/97, Kibbey *et al.*, Method and System for Identification, Purification and Quantitation of Reaction Components;
- EP0617048, 1994-09-28, WATERS INVESTMENTS LTD (US), Method of Capillary Isoelectric Focusing of Proteins and Peptides with Fraction Collection for Post-Run

Analysis;

- WO9840395/US9804875/GB9703307, 1998-09-17, TAYLOR *et al.*, Band Array Display of Polynucleotide Separations; and
- WO9701755, 1997-01-16, PERSEPTIVE BIOSYSTEMS INC (US), High Speed, Automated, Continuous Flow, Multi-Dimensional Molecular Selection and Analysis.
- U.S. 5131998, Jorgenson, 07/21/92, Two-Dimensional High-Performance Liquid Chromatography/Capillary Electrophoresis;
- WO 98/23950 describes computer assisted methods and apparatus for identifying, selecting, and characterizing molecules in a biological sample. The method generates a two-dimensional array; and
- Toribio *et al.*, J. Chromat. B 684:1 [1996] reviews methods for the chromatographic and electrophoretic purification of pyridine nucleotide-dependent enzymes.

Applicants have become aware of the following printed publications which may be material to the examination of this application:

- Houen *et al.*, Characterization of protein carboxy-terminal ends using carboxypeptidase peptide Y: Sequence, composition, and identification of the carboxy-terminal peptide by peptide mapping, Methods Molecular Cellular Biology, 3(4):175-182 (1992) (abstract only);
- Raznukov *et al.*, Selective digital filtering of mass spectra of chromatography data for determination of "target" compounds in complex mixtures, Advances in Mass Spectrometry, 14 (EO44280/44281-EO44280/44211) (1998) (abstract only);
- SZE *et al.*, Time-of-flight effects in matrix-assisted laser desorption/ionization Fourier transform mass spectrometry, Rapid Commun Mass Spectrom, 13(5):398-406 (1999) (abstract only);
- Wall *et al.*, Isoelectric focusing nonporous silica reversed-phase high-performance

liquid chromatography/electrospray ionization time-of-flight mass spectrometry: a three-dimensional liquid-phase protein separation method as applied to the human erythroleukemia cell-line, *Rapid Commun Mass Spectrom*, 15(18):1649-1661 (2001);

- Wall *et al.*, Three-dimensional protein map according to pI, hydrophobicity and molecular mass, *J Chromatogr B Analyt Technol Biomed Life Sci*, 774(1):53-58 (2002); and
- Zgoda *et al.*, Prediction and experimental confirmation of the cytochrome b5 three-dimensional peptide map, *Physical Chemical Biology and Medicine*, 2(3):135-142 (1995) (abstract only)
- Wilkins and Williams, Cross-species protein identification using amino acid composition, peptide mass fingerprinting, isoelectric point and molecular mass: a theoretical evaluation, *J Theor Biol*, 186(1):7-15 (1997)
- Anderson *et al.*, A two-dimensional gel database of rat liver proteins useful in gene regulation and drug effects studies, *Electrophoresis*, 12(11):907-930 (1991);
- Damerval, Quantification of silver-stained proteins resolved by two-dimensional electrophoresis: genetic variability as related to abundance and solubility in two maize lines, *Electrophoresis*, 15(12):1573-1579 (1994);
- Kahn, From genome to proteome: looking at a cell's proteins, *Science*, 270(5235):369-370 (1995);
- Neidhardt *et al.*, Genomically linked cellular protein databases derived from two-dimensional polyacrylamide gel electrophoresis, *Electrophoresis*, 10(2):116-122 (1989);
- O'Farrell, High resolution two-dimensional electrophoresis of proteins, *J Biol Chem*, 250(10):4007-4021 (1975);
- Patterson, Matrix-assisted laser-desorption/ionization mass spectrometric approaches for the identification of gel-separated proteins in the 5-50 pmol range, *Electrophoresis*, 16(7):1104-1114 (1995);
- Aebersold *et al.*, Internal amino acid sequence analysis of proteins separated by one-

or two-dimensional gel electrophoresis after in situ protease digestion on nitrocellulose, Proc Natl Acad Sci U S A, 84(20):6970-6974 (1987);

- Brown and Cooper, Regulation, substrates and functions of src, Biochim Biophys Acta, 1287(2-3):121-149 (1996);
- Cohen and Chait, Mass spectrometry of whole proteins eluted from sodium dodecyl sulfate-polyacrylamide gel electrophoresis gels, Anal Biochem, 247(2):257-267 (1997);
- Courchesne *et al.*, Comparison of in-gel and on-membrane digestion methods at low to sub-pmol level for subsequent peptide and fragment-ion mass analysis using matrix-assisted laser-desorption/ionization mass spectrometry, Electrophoresis, 18(3-4):369-381 (1997);
- Davidsson *et al.*, Characterization of proteins from human cerebrospinal fluid by a combination of preparative two-dimensional liquid-phase electrophoresis and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, Anal Chem, 71(3):642-647 (1999);
- Dawson *et al.*, MCF10AT: a model for the evolution of cancer from proliferative breast disease, Am J Pathol, 148(1):313-319 (1996);
- Egan *et al.*, Activation of Src in human breast tumor cell lines: elevated levels of phosphotyrosine phosphatase activity that preferentially recognizes the Src carboxy terminal negative regulatory tyrosine 530, Oncogene, 18(5):1227-1237 (1999);
- Gottleib and Oren, p53 in growth control and neoplasia, Biochimica et Biophysica Acta, 1287(2-3):77-102 (1996). This publication describes a structural and functional review of p53;
- Hayakawa *et al.*, Isoelectric focusing of biotinidase and lipoamidase with the addition of non-ionic detergent, Anal Chim Acta, 372(1-2):281-289 (1998);
- Haynes *et al.*, Proteome analysis: biological assay or data archive?, Electrophoresis, 19(11):1862-1871 (1998) (abstract only);
- Henzel *et al.*, Identifying proteins from two-dimensional gels by molecular mass

searching of peptide fragments in protein sequence databases, Proc Natl Acad Sci U S A, 90(11):5011-5015 (1993);

- Lahm and Langen, Mass spectrometry: a tool for the identification of proteins separated by gels, 21(2):411-429 (2000);
- Li *et al.*, Development of a three-dimensional topographic map display for capillary electrophoresis/mass spectrometry with an ion trap/reflectron time-of-flight mass spectrometer detector: applications to tryptic digests of isoforms of myelin basic protein, J Am Soc Mass Spectrom, 9(7):701-709 (1998);
- Liang *et al.*, Characterization of SDS--PAGE-separated proteins by matrix-assisted laser desorption/ionization mass spectrometry, Anal Chem, 68(6):1012-1018 (1996);
- Liang and Pardee, Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction, Science, 257(5072):967-971 (1992);
- Link *et al.*, Direct analysis of protein complexes using mass spectrometry, Nat Biotechnol, 17(7):676-682 (1999);
- Loo *et al.*, Mass Spectrometry of Proteins Directly from Polyacrylamide Gels, Anal Chem, 68(11):1910 - 1917 (1996);
- Lopez, Better approaches to finding the needle in a haystack: optimizing proteome analysis through automation, Electrophoresis, 21(6):1082-1093 (2000);
- MacNair *et al.*, Rapid separation and characterization of protein and peptide mixtures using 1.5 microns diameter non-porous silica in packed capillary liquid chromatography/mass spectrometry, Rapid Commun Mass Spectrom, 11(12):1279-1285 (1997);
- Maggiolini *et al.*, Adrenal androgens stimulate the proliferation of breast cancer cells as direct activators of estrogen receptor alpha, Cancer Res, 59(19):4864-4869 (1999);
- Mao *et al.*, Activation of c-Src by receptor tyrosine kinases in human colon cancer cells with high metastatic potential, Oncogene, 15(25):3083-3090 (1997);
- Michael, Detection of electrospray ionization using a quadrupole ion trap

storage/reflectron time-of-flight mass spectrometer, *Anal Chem*, 65(2614-2620 (1994);

- Miller *et al.*, Differential display, subtractive hybridization, and application of methodology to search for point mutations to identify genetic defects responsible for progression in MCF10AT model of human breast disease, *Electrophoresis*, 20(2):256-260 (1999);

- Miller *et al.*, Xenograft model of progressive human proliferative breast disease, *J Natl Cancer Inst*, 85(21):1725-1732 (1993);

- Mills *et al.*, Prospective study of exogenous hormone use and breast cancer in Seventh-day Adventists, *Cancer*, 64(3):591-597 (1989);

- Molinari *et al.*, Estradiol induces functional inactivation of p53 by intracellular redistribution, *Cancer Res*, 60(10):2594-2597 (2000);

- Nilsson *et al.*, Identification of proteins in a human pleural exudate using two-dimensional preparative liquid-phase electrophoresis and matrix-assisted laser desorption/ionization mass spectrometry, *Electrophoresis*, 20(4-5):860-865 (1999);

- Ottenhoff-Kalff *et al.*, Characterization of protein tyrosine kinases from human breast cancer: involvement of the c-src oncogene product, *Cancer Res*, 52(17):4773-4778 (1992);

- Ozturk *et al.*, Detection of c-erbB-2 mRNAs using dig-labelled oligonucleotide probe with in situ hybridisation in human breast carcinoma: comparison with immunohistochemical results, *Anal Cell Pathol*, 16(4):201-209 (1998);

- Porter *et al.*, Role of estrogen receptor/Sp1 complexes in estrogen-induced heat shock protein 27 gene expression, *Mol Endocrinol*, 10(11):1371-1378 (1996);

- Qian and Lubman, Analysis of tryptic digests using microbore HPLC with an ion trap storage/reflectron time-of-flight detector, *Anal Chem*, 67(17):2870-2877 (1995);

- Rodriguez *et al.*, Towards stoichiometric silver staining of proteins resolved in complex two-dimensional electrophoresis gels: real-time analysis of pattern development, *Electrophoresis*, 14(7):628-637 (1993);

- Rosen *et al.*, Analysis of pp60c-src protein kinase activity in human tumor cell lines and tissues, J Biol Chem, 261(29):13754-13759 (1986);
- Ruddon, Cancer biology, Oxford University Press, 3rd, 318-340 (1995);
- Russell *et al.*, Cyclin D1 and D3 associate with the SCF complex and are coordinately elevated in breast cancer, Oncogene, 18(11):1983-1991 (1999);
- Santner *et al.*, MCF-10CA1 cell lines: New highly tumorigenic derivatives of the MCF-10AT system, Proc Am Assoc Cancer Res, 39(202 (1998) (abstract only);
- Schweinfest and Pappas, Subtraction hybridization: an approach to the isolation of genes differentially expressed in cancer and other biological system (Review), Intern J Oncol, 1(499-506 (1992);
- Shekhar *et al.*, Transcriptional activation of functional endogenous estrogen receptor gene expression in MCF10AT cells: a model for early breast cancer, Int J Oncol, 13(5):907-915 (1998);
- Shevchenko *et al.*, Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels, Anal Chem, 68(5):850-858 (1996);
- Soule *et al.*, Isolation and characterization of a spontaneously immortalized human breast epithelial cell line, MCF-10, Cancer Res, 50(18):6075-6086 (1990);
- Sturtevant, Applications of differential-display reverse transcription-PCR to molecular pathogenesis and medical mycology, Clin Microbiol Rev, 13(3):408-427 (2000);
- Tesarik *et al.*, Estradiol modulates breast cancer cell apoptosis: a novel nongenomic steroid action relevant to carcinogenesis, Steroids, 64(1-2):22-27 (1999);
- Tetu *et al.*, Prognostic significance of heat-shock protein-27 in node-positive breast carcinoma: an immunohistochemical study, Breast Cancer Res Treat, 36(1):93-97 (1995);
- Thomas *et al.*, Exogenous estrogens and other risk factors for breast cancer in women with benign breast diseases, J Natl Cancer Inst, 69(5):1017-1025 (1982);

- Williams *et al.*, Analysis of differential protein expression in normal and neoplastic human breast epithelial cell lines, *Electrophoresis*, 19(2):333-343 (1998);
- Yang and Lee, Capillary isoelectric focusing-electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry for protein characterization, *Anal Chem*, 70(15):3235-3241 (1998);
- Figeys and Aebersold, Nanoflow solvent gradient delivery from a microfabricated device for protein identifications by electrospray ionization mass spectrometry, *Anal Chem*, 70(18):3721-3727 (1998) (abstract only);
- Figeys *et al.*, Data-dependent modulation of solid-phase extraction capillary electrophoresis for the analysis of complex peptide and phosphopeptide mixtures by tandem mass spectrometry: application to endothelial nitric oxide synthase, *Anal Chem*, 71(13):2279-2287 (1999) (abstract only);
- Figeys *et al.*, Protein identification by solid phase microextraction-capillary zone electrophoresis-microelectrospray-tandem mass spectrometry, *Nat Biotechnol*, 14(11):1579-1583 (1996) (abstract only);
- Figeys *et al.*, "An integrated microfluidics-tandem mass spectrometry system for automated protein analysis," *Anal. Chem.* 70:3728 (1998) (Abstract only);
- Figeys *et al.*, Microfabricated device coupled with an electrospray ionization quadrupole time-of-flight mass spectrometer: protein identifications based on enhanced-resolution mass spectrometry and tandem mass spectrometry data, *Rapid Commun Mass Spectrom*, 12(20):1435-1444 (1998) (Abstract only);
- Figeys *et al.*, A microfabricated device for rapid protein identification by microelectrospray ion trap mass spectrometry, *Anal Chem*, 69(16):3153-3160 (1997);
- Figeys *et al.*, Optimization of solid phase microextraction - capillary zone electrophoresis - mass spectrometry for high sensitivity protein identification, *Electrophoresis*, 19(13):2338-2347 (1998) (abstract only);

- Hunt *et al.*, Characterization of peptides bound to the class I MHC molecule HLA-A2.1 by mass spectrometry, *Science*. 1992 Mar 6;255(5049):1261-3 (abstract only);
- Henderson *et al.*, HLA-A2.1-associated peptides from a mutant cell line: a second pathway of antigen presentation, *Science*. 1992 Mar 6;255(5049):1264-6 (abstract only);
- Rossomando *et al.*, Identification of Tyr-185 as the site of tyrosine autophosphorylation of recombinant mitogen-activated protein kinase p42mapk, *Proc Natl Acad Sci U S A*. 1992 Jul 1;89(13):5779-83;
- Henderson *et al.*, Direct identification of an endogenous peptide recognized by multiple HLA-A2.1-specific cytotoxic T cells, *Proc Natl Acad Sci U S A*, 90(21): 10275-9 (1993);
- Kingan *et al.*, The loss of female sex pheromone after mating in the corn earworm moth *Helicoverpa zea*: identification of a male pheromonostatic peptide. *Proc Natl Acad Sci U S A*. 1995 May 23;92(11):5082-6;
- Hungerford *et al.*, Identification of a Novel Marker for Primordial Smooth Muscle and Its Differential Expression Pattern in Contractile vs Noncontractile Cells, *J. Cell Biol.* 137(4) May 19, 1997: 925-937;
- Pierce *et al.*, Cutting Edge: The HLA-A*0101-Restricted HY Minor Histocompatibility Antigen Originates from DFFRY and Contains a Cysteinylation Cysteine Residue as Identified by a Novel Mass Spectrometric Technique, *The Journal of Immunology*, 163: 6360-6364 (1999);
- Martin *et al.*, Subfemtomole MS and MS/MS Peptide Sequence Analysis Using Nano-HPLC Micro-ESI Fourier Transform Ion Cyclotron Resonance Mass Spectrometry, *Anal. Chem.*; 72(18) pp 4266 - 4274 (2000) (abstract only);
- Flyer, *et al.*, Identification by Mass Spectrometry of CD8+-T-Cell Mycobacterium tuberculosis Epitopes within the Rv0341 Gene Product, *Infection and Immunity*, 70(6):2926-2932, (June 2002);

- U.S. 6,139,734, Settlage, *et al.*;
- U. S. 6,315,905, Settlage, *et al.*;
- United States Patent Application 20030143647 A1, Finn, , *et al.*;

This Information Disclosure Statement under 37 C.F.R. §§ 1.56 and 1.97 is not to be construed as a representation that a search has been made, that additional information material to the examination of this application does not exist, or that any one or more of these citations constitutes prior art.

Dated: October 10, 2003



Tanya A. Arenson

Registration No. 47,391

MEDLEN & CARROLL, LLP

101 Howard Street, Suite 350

San Francisco, California 94105

415/904-6500

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket No.: UM-06102

Serial No.: 09/778,548

INFORMATIONAL DISCLOSURE STATEMENT BY APPLICANT

(Several Sheets If Necessary)

Applicant: David M. Lubman *et al.*

(37 CFR § 1.98(b))

Filing Date: 02/07/01

Group Art Unit:

U.S. PATENT DOCUMENTS

Examiner Initials	Cite No.	Serial / Patent Number	Issue Date	Applicant / Patentee	Class	Subclass	Filing Date
	1	6,002,127	12/14/99	Vestal			
	2	5,498,545	3/12/96	Vesta			
	3	5,670,054	9/23/97	Kibbey <i>et al.</i>			
	4	5,131,998	7/21/92	Jorgenson			
	5	6,139,734		Settlage <i>et al.</i>			
	6	6,315,905		Settlage <i>et al.</i>			
	7	20030143647 A1					
	8	0020039747	02/07/01	Lubman, <i>et al.</i>			
	9	20030064527	04/26/02	Lubman, <i>et al.</i>			
	10	20030058239	04/26/02,	Wall, <i>et al.</i>			
	11	09/968930	10/1/01,	Lubman, <i>et al.</i>			
	12	10/448,564	5/30/03,	Lubman, <i>et al.</i>			

FOREIGN PATENTS OR PUBLISHED FOREIGN PATENT APPLICATIONS

		Document Number	Publication Date	Country / Patent Office	Class	Subclass	Translation	
							Yes	No
	13	WO 01/58925	08/16/01	PCT				
	14	WO 02/088701	11/07/02	PCT				
	15	WO 01/58926	03/2/03	PCT				
	16	WO 01/59460	03/2/03	PCT				
	17	PCT/US03/16899	5/30/03	PCT				
	18	EP 0617048	9/28/94	EP				
	19	WO 98/40395	9/17/98	PCT				
	20	WO 98/23950		PCT				
	21	WO 97/01755	1/16/97	PCT				

OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

22	Rossomando <i>et al.</i> , Identification of Tyr-185 as the site of tyrosine autophosphorylation of recombinant mitogen-activated protein kinase p42mapk, Proc Natl Acad Sci U S A. 1992 Jul 1;89(13):5779-83;
23	Henderson <i>et al.</i> , Direct identification of an endogenous peptide recognized by multiple HLA-A2.1-specific cytotoxic T cells, Proc Natl Acad Sci U S A, 90(21): 10275-9 (1993);
24	Kingan <i>et al.</i> , The loss of female sex pheromone after mating in the corn earworm moth <i>Helicoverpa zea</i> : identification of a male pheromonostatic peptide. Proc Natl Acad Sci U S A. 1995 May 23;92(11):5082-6;
25	Hungerford <i>et al.</i> , Identification of a Novel Marker for Primordial Smooth Muscle and Its Differential Expression Pattern in Contractile vs Noncontractile Cells, J. Cell Biol. 137(4) May 19, 1997: 925-937;
26	Pierce <i>et al.</i> , Cutting Edge: The HLA-A*0101-Restricted HY Minor Histocompatibility Antigen Originates from DFFRY and Contains a Cysteinylylated Cysteine Residue as Identified by a Novel Mass Spectrometric Technique, The Journal of Immunology, 163: 6360-6364 (1999);
27	Martin <i>et al.</i> , Subfemtomole MS and MS/MS Peptide Sequence Analysis Using Nano-HPLC Micro-ESI Fourier Transform Ion Cyclotron Resonance Mass Spectrometry, Anal. Chem.; 72(18) pp 4266 - 4274 (2000) (abstract only);

EXAMINER:

Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket No.: UM-06102

Serial No.: 09/778,548

INFORMATION DISCLOSURE STATEMENT BY APPLICANT

(Use Several Sheets If Necessary)

Applicant: David M. Lubman *et al.*

Filing Date: 02/07/01

Group Art Unit:

(37 CFR § 1.98(b))

OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

28	Ayala <i>et al.</i> , Use of Rotofor preparative isoelectrofocusing cell in protein purification procedure, Appl Biochem Biotechnol, 69(1):11-16 (1998)
29	Bini <i>et al.</i> , Protein expression profiles in human breast ductal carcinoma and histologically normal tissue, Electrophoresis, 18(15):2832-2841 (1997)
30	Chen <i>et al.</i> , Identification of proteins from two-dimensional gel electrophoresis of human erythroleukemia cells using capillary high performance liquid chromatography/electrospray-ion trap-reflectron time-of-flight mass spectrometry with two-dimensional topographic map analysis of in-gel tryptic digest products, Rapid Commun Mass Spectrom, 13(19):1907-1916 (1999)
31	Chen <i>et al.</i> , Rapid identification and screening of proteins from whole cell lysates of human erythroleukemia cells in the liquid phase, using non-porous reversed phase high-performance liquid chromatography separations of proteins followed by matrix-assisted [correction of multi-assisted] laser desorption/ionization mass spectrometry analysis and sequence database searching, Rapid Commun Mass Spectrom, 12(24):1994-2003 (1998)
32	Chong <i>et al.</i> , Rapid screening of protein profiles of human breast cancer cell lines using non-porous reversed-phase high performance liquid chromatography separation with matrix-assisted laser desorption/ionization time-of-flight mass spectral analysis, Rapid Commun Mass Spectrom, 13(18):1808-1812 (1999)
33	Dai <i>et al.</i> , Two-layer sample preparation: a method for MALDI-MS analysis of complex peptide and protein mixtures, Anal Chem, 71(5):1087-1091 (1999)
34	Figeys and Aebersold, Microfabricated modules for sample handling, sample concentration and flow mixing: application to protein analysis by tandem mass spectrometry, J Biomech Eng, 121(1):7-12 (1999)
35	Fuqua <i>et al.</i> , Induction of the estrogen-regulated "24K" protein by heat shock, Cancer Res, 49(15):4126-4129 (1989)
36	Griffin <i>et al.</i> , Direct database searching with MALDI-PSD spectra of peptides, Rapid Commun Mass Spectrom, 9(15):1546-1551 (1995)
37	Hanash, Advances in Electrophoresis, 1-44, (1998)
38	Herbert, Advances in protein solubilisation for two-dimensional electrophoresis, Electrophoresis, 20(4-5):660-663 (1999)
39	Immler <i>et al.</i> , Identification of phosphorylated proteins from thrombin-activated human platelets isolated by two-dimensional gel electrophoresis by electrospray ionization-tandem mass spectrometry (ESI-MS/MS) and liquid chromatography-electrospray ionization-mass spectrometry (LC-ESI-MS), Electrophoresis, 19(6):1015-1023 (1998)
40	Liang <i>et al.</i> , Determination of bacterial protein profiles by matrix-assisted laser desorption/ionization mass spectrometry with high-performance liquid chromatography, Rapid Commun Mass Spectrom, 10(10):1219-1226 (1996)
41	Matsui <i>et al.</i> , Immobilized pH gradient two-dimensional gel electrophoresis and mass spectrometric identification of cytokine-regulated proteins in ME-180 cervical carcinoma cells, Electrophoresis, 18(3-4):409-417 (1997)
42	Mohammad <i>et al.</i> , Induced expression of alpha-enolase in differentiated diffuse large cell lymphoma, Enzyme Protein, 48(1):37-44 (1994)
43	Neubauer and Mann, Mapping of phosphorylation sites of gel-isolated proteins by nanoelectrospray tandem mass spectrometry: potentials and limitations, Anal Chem, 71(1):235-242 (1999)
44	Nilsson <i>et al.</i> , Isolation and characterization of proteins from human lymphocyte nuclei using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry and post-source decay analysis, Rapid Commun Mass Spectrom, 11(6):610-612 (1997)
45	Opitck <i>et al.</i> , Comprehensive two-dimensional high-performance liquid chromatography for the isolation of overexpressed proteins and proteome mapping, Anal Biochem, 258(2):349-361 (1998)
46	Patterson and Aebersold, Mass spectrometric approaches for the identification of gel-separated proteins, Electrophoresis, 16(10):1791-1814 (1995)
47	Pinto <i>et al.</i> , An enhanced microfluidic chip coupled to an electrospray Qstar mass spectrometer for protein identification, Electrophoresis, 21(1):181-190 (2000)
48	Rasmussen <i>et al.</i> , Two-dimensional gel database of human breast carcinoma cell expressed proteins: an update, Electrophoresis, 19(5):818-825 (1998)
49	Redner <i>et al.</i> , The t(5;17) variant of acute promyelocytic leukemia expresses a nucleophosmin-retinoic acid receptor fusion, Blood, 87(3):882-886 (1996)

Examiner:

Date Considered:

EXAMINER:

Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

RECEIVED
 OCT 17 2003
 RECH CENTER 1601/2900

FORM PTO-1449 (Modified)		U.S. Department of Commerce Patent and Trademark Office		Attorney Docket No.: UM-06102	Serial No.: 09/778,548
INFORMATION DISCLOSURE STATEMENT BY APPLICANT (Use Several Sheets If Necessary)				Applicant: David M. Lubman <i>et al.</i>	
				Filing Date: 02/07/01	Group Art Unit:
(37 CFR § 1.98(b))					
OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)					
	50	Reid <i>et al.</i> , Capillary column chromatography improves sample preparation for mass spectrometric analysis: complete characterization of human alpha-enolase from two-dimensional gels following in situ proteolytic digestion, <i>Electrophoresis</i> , 19(6):946-955 (1998)			
	51	Righetti, Isoelectric focusing : theory, methodology, and applications, Elsevier Biomedical Press; Elsevier-North Holland, 1st, (1983)			
	52	Rosenfeld <i>et al.</i> , In-gel digestion of proteins for internal sequence analysis after one- or two-dimensional gel electrophoresis, <i>Anal Biochem</i> , 203(1):173-179 (1992)			
	53	Sanchez <i>et al.</i> , Inside SWISS-2DPAGE database, <i>Electrophoresis</i> , 16(7):1131-1151 (1995)			
	54	Sirover, New insights into an old protein: the functional diversity of mammalian glyceraldehyde-3-phosphate dehydrogenase, <i>Biochim Biophys Acta (BBA) - Protein Structure and Molecular Enzymology</i> , 1432(2):159-184 (1999)			
	55	Steller, Mechanisms and genes of cellular suicide, <i>Science</i> , 267(5203):1445-1449 (1995)			
	56	ten Hoeve <i>et al.</i> , Isolation and chromosomal localization of CRKL, a human crk-like gene, <i>Oncogene</i> , 8(9):2469-2474 (1993)			
	57	Wall <i>et al.</i> , Isoelectric focusing nonporous RP HPLC: a two-dimensional liquid-phase separation method for mapping of cellular proteins with identification using MALDI-TOF mass spectrometry, <i>Anal Chem</i> , 72(6):1099-1111 (2000)			
	58	Wall <i>et al.</i> , Rapid profiling of induced proteins in bacteria using MALDI-TOF mass spectrometric detection of nonporous RP HPLC-separated whole cell lysates, <i>Anal Chem</i> , 71(17):3894-3900 (1999)			
	59	Welsh <i>et al.</i> , Variation in expression of hsp27 messenger ribonucleic acid during the cycle of the seminiferous epithelium and co-localization of hsp27 and microfilaments in Sertoli cells of the rat, <i>Biol Reprod</i> , 55(1):141-151 (1996)			
	60	Yates, Mass spectrometry and the age of the proteome, <i>J Mass Spectrom</i> , 33(1):1-19 (1998)			
	61	Zugaro <i>et al.</i> , Characterization of rat brain stathmin isoforms by two-dimensional gel electrophoresis-matrix assisted laser desorption/ionization and electrospray ionization-ion trap mass spectrometry, <i>Electrophoresis</i> , 19(5):867-876 (1998)			
	62	Chong <i>et al.</i> , Rapid profiling of E. coli proteins up to 500 kDa from whole cell lysates using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, <i>Rapid Commun Mass Spectrom</i> , 11(17):1900-1908 (1997)			
	63	Opitck <i>et al.</i> , Two-dimensional SEC/RPLC coupled to mass spectrometry for the analysis of peptides, <i>Anal Chem</i> , 69(13):2283-2291 (1997)			
	64	Regnier and Huang, Future potential of targeted component analysis by multidimensional liquid chromatography-mass spectrometry, <i>J Chromatogr A</i> , 750(1-2):3-10 (1996)			
	65	Andrews <i>et al.</i> , Analysis of DNA adducts using high-performance separation techniques coupled to electrospray ionization mass spectrometry, <i>J Chromatogr A</i> , 856(1-2):515-526 (1999)			
	66	Ball, Purification of synthetic peptides with the aid of reversible chromatographic probes, <i>J Chromatogr A</i> , 686(73-83 (1994)			
	67	Binz <i>et al.</i> , A molecular scanner to automate proteomic research and to display proteome images, <i>Anal Chem</i> , 71(21):4981-4988 (1999)			
	68	Davidsson and Nilsson, Peptide mapping of proteins in cerebrospinal fluid utilizing a rapid preparative two-dimensional electrophoretic procedure and matrix-assisted laser desorption/ionization mass spectrometry, <i>Biochim Biophys Acta</i> , 1473(2-3):391-399 (1999)			
	69	Lee, Protein separation using non-porous sorbents, <i>J Chromatogr B Biomed Sci Appl</i> , 699(1-2):29-45 (1997)			
	70	Medzihradsky <i>et al.</i> , Protein sequence and structural studies employing matrix-assisted laser desorption ionization-high energy collision-induced dissociation, <i>International Journal of Mass Spectrometry and Ion Processes</i> , 160(1-3):357-369 (1997)			
	71	Nimura <i>et al.</i> , Fast protein separation by reversed-phase high-performance liquid chromatography on octadecylsilyl-bonded non-porous silica gel. Effect of particle size of column packing on column efficiency., <i>J Chromatogr</i> , 585(207-211 (1991)			
	72	Richmond, High-throughput flow injection analysis-mass spectrometry with networked delivery of colour rendered results: the characterisation of liquid chromatography fractions, <i>J Chromatogr A</i> , 835(29-39 (1999)			
	73	Wilkins <i>et al.</i> , From proteins to proteomes: large scale protein identification by two-dimensional electrophoresis and amino acid analysis, <i>Biotechnology (N Y)</i> , 14(1):61-65 (1996)			
Examiner:			Date Considered:		
EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.					

RECEIVED
 OCT 17 2003
 TECH CENTER 160012900

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket No.: UM-06102

Serial No.: 09/778,548

INFORMATION DISCLOSURE STATEMENT BY APPLICANT

(Use General Sheets If Necessary)

Applicant: David M. Lubman *et al.*

Filing Date: 02/07/01

Group Art Unit:

(37 CFR § 1.98(b))


OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

74	Toribio <i>et al.</i> , J. Chromat. B 684:1 [1996] reviews methods for the chromatographic and electrophoretic purification of pyridine nucleotide-dependent enzymes
75	Houen <i>et al.</i> , Characterization of protein carboxy-terminal ends using carboxypeptidase peptide Y: Sequence, composition, and identification of the carboxy-terminal peptide by peptide mapping, Methods Molecular Cellular Biology, 3(4):175-182 (1992) (abstract only)
76	Raznikov <i>et al.</i> , Selective digital filtering of mass spectra of chromatography data for determination of "target" compounds in complex mixtures, Advances in Mass Spectrometry, 14 (EO44280/44281-EO44280/44211) (1998) (abstract only)
77	SZE <i>et al.</i> , Time-of-flight effects in matrix-assisted laser desorption/ionization Fourier transform mass spectrometry, Rapid Commun Mass Spectrom, 13(5):398-406 (1999) (abstract only)
78	Wall <i>et al.</i> , Isoelectric focusing nonporous silica reversed-phase high-performance liquid chromatography/electrospray ionization time-of-flight mass spectrometry: a three-dimensional liquid-phase protein separation method as applied to the human erythroleukemia cell-line, Rapid Commun Mass Spectrom, 15(18):1649-1661 (2001)
79	Wall <i>et al.</i> , Three-dimensional protein map according to pI, hydrophobicity and molecular mass, J Chromatogr B Analyt Technol Biomed Life Sci, 774(1):53-58 (2002)
80	Zgoda <i>et al.</i> , Prediction and experimental confirmation of the cytochrome b5 three-dimensional peptide map, Physical Chemical Biology and Medicine, 2(3):135-142 (1995) (abstract only)
81	Wilkins and Williams, Cross-species protein identification using amino acid composition, peptide mass fingerprinting, isoelectric point and molecular mass: a theoretical evaluation, J Theor Biol, 186(1):7-15 (1997) (abstract only)
82	Anderson <i>et al.</i> , A two-dimensional gel database of rat liver proteins useful in gene regulation and drug effects studies, Electrophoresis, 12(11):907-930 (1991)
83	Damerval, Quantification of silver-stained proteins resolved by two-dimensional electrophoresis: genetic variability as related to abundance and solubility in two maize lines, Electrophoresis, 15(12):1573-1579 (1994)
84	Kahn, From genome to proteome: looking at a cell's proteins, Science, 270(5235):369-370 (1995);
85	Neidhardt <i>et al.</i> , Genomically linked cellular protein databases derived from two-dimensional polyacrylamide gel electrophoresis, Electrophoresis, 10(2):116-122 (1989)
86	O'Farrell, High resolution two-dimensional electrophoresis of proteins, J Biol Chem, 250(10):4007-4021 (1975)
87	Patterson, Matrix-assisted laser-desorption/ionization mass spectrometric approaches for the identification of gel-separated proteins in the 5-50 pmol range, Electrophoresis, 16(7):1104-1114 (1995)
88	Aebersold <i>et al.</i> , Internal amino acid sequence analysis of proteins separated by one- or two-dimensional gel electrophoresis after in situ protease digestion on nitrocellulose, Proc Natl Acad Sci U S A, 84(20):6970-6974 (1987)
89	Brown and Cooper, Regulation, substrates and functions of src, Biochim Biophys Acta, 1287(2-3):121-149 (1996)
90	Cohen and Chait, Mass spectrometry of whole proteins eluted from sodium dodecyl sulfate-polyacrylamide gel electrophoresis gels, Anal Biochem, 247(2):257-267 (1997)
91	Courchesne <i>et al.</i> , Comparison of in-gel and on-membrane digestion methods at low to sub-pmol level for subsequent peptide and fragment-ion mass analysis using matrix-assisted laser-desorption/ionization mass spectrometry, Electrophoresis, 18(3-4):369-381 (1997)
92	Davidsson <i>et al.</i> , Characterization of proteins from human cerebrospinal fluid by a combination of preparative two-dimensional liquid-phase electrophoresis and matrix-assisted laser desorption/ionization time-of-flight mass spectrometry, Anal Chem, 71(3):642-647 (1999)
93	Dawson <i>et al.</i> , MCF10AT: a model for the evolution of cancer from proliferative breast disease, Am J Pathol, 148(1):313-319 (1996)
94	Egan <i>et al.</i> , Activation of Src in human breast tumor cell lines: elevated levels of phosphotyrosine phosphatase activity that preferentially recognizes the Src carboxy terminal negative regulatory tyrosine 530, Oncogene, 18(5):1227-1237 (1999)
95	Gottlieb and Oren, p53 in growth control and neoplasia, Biochimica et Biophysica Acta, 1287(2-3):77-102 (1996). This publication describes a structural and functional review of p53
96	Hayakawa <i>et al.</i> , Isoelectric focusing of biotinidase and lipoamidase with the addition of non-ionic detergent, Anal Chim Acta, 372(1-2):281-289 (1998)
97	Haynes <i>et al.</i> , Proteome analysis: biological assay or data archive?, Electrophoresis, 19(11):1862-1871 (1998) (abstract only)

Examiner:

Date Considered:

EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.



RECEIVED
OCT 14 2003
TECH CENTER 1100 2900

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket No.: UM-06102

Serial No.: 09/778,548

INFORMATION ENCLOSURE STATEMENT BY APPLICANT

(Use Several Sheets if Necessary)

Applicant: David M. Lubman *et al.*

Filing Date: 02/07/01

Group Art Unit:

(37 CFR § 1.98(b))

OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

98	Henzel <i>et al.</i> , Identifying proteins from two-dimensional gels by molecular mass searching of peptide fragments in protein sequence databases, <i>Proc Natl Acad Sci U S A</i> , 90(11):5011-5015 (1993)
99	Lahm and Langen, Mass spectrometry: a tool for the identification of proteins separated by gels, 21(2):411-429 (2000)
100	Li <i>et al.</i> , Development of a three-dimensional topographic map display for capillary electrophoresis/mass spectrometry with an ion trap/reflectron time-of-flight mass spectrometer detector: applications to tryptic digests of isoforms of myelin basic protein, <i>J Am Soc Mass Spectrom</i> , 9(7):701-709 (1998)
101	Liang <i>et al.</i> , Characterization of SDS--PAGE-separated proteins by matrix-assisted laser desorption/ionization mass spectrometry, <i>Anal Chem</i> , 68(6):1012-1018 (1996)
102	Liang and Pardee, Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction, <i>Science</i> , 257(5072):967-971 (1992)
103	Link <i>et al.</i> , Direct analysis of protein complexes using mass spectrometry, <i>Nat Biotechnol</i> , 17(7):676-682 (1999)
104	Loo <i>et al.</i> , Mass Spectrometry of Proteins Directly from Polyacrylamide Gels, <i>Anal Chem</i> , 68(11):1910 - 1917 (1996)
105	Lopez, Better approaches to finding the needle in a haystack: optimizing proteome analysis through automation, <i>Electrophoresis</i> , 21(6):1082-1093 (2000)
106	MacNair <i>et al.</i> , Rapid separation and characterization of protein and peptide mixtures using 1.5 microns diameter non-porous silica in packed capillary liquid chromatography/mass spectrometry, <i>Rapid Commun Mass Spectrom</i> , 11(12):1279-1285 (1997)
107	Maggiolini <i>et al.</i> , Adrenal androgens stimulate the proliferation of breast cancer cells as direct activators of estrogen receptor alpha, <i>Cancer Res</i> , 59(19):4864-4869 (1999)
108	Mao <i>et al.</i> , Activation of c-Src by receptor tyrosine kinases in human colon cancer cells with high metastatic potential, <i>Oncogene</i> , 15(25):3083-3090 (1997)
109	Michael, Detection of electrospray ionization using a quadrupole ion trap storage/reflectron time-of-flight mass spectrometer, <i>Anal Chem</i> , 65(2614-2620 (1994)
110	Miller <i>et al.</i> , Differential display, subtractive hybridization, and application of methodology to search for point mutations to identify genetic defects responsible for progression in MCF10AT model of human breast disease, <i>Electrophoresis</i> , 20(2):256-260 (1999)
111	Miller <i>et al.</i> , Xenograft model of progressive human proliferative breast disease, <i>J Natl Cancer Inst</i> , 85(21):1725-1732 (1993)
112	Mills <i>et al.</i> , Prospective study of exogenous hormone use and breast cancer in Seventh-day Adventists, <i>Cancer</i> , 64(3):591-597 (1989)
113	Molinari <i>et al.</i> , Estradiol induces functional inactivation of p53 by intracellular redistribution, <i>Cancer Res</i> , 60(10):2594-2597 (2000)
114	Nilsson <i>et al.</i> , Identification of proteins in a human pleural exudate using two-dimensional preparative liquid-phase electrophoresis and matrix-assisted laser desorption/ionization mass spectrometry, <i>Electrophoresis</i> , 20(4-5):860-865 (1999)
115	Ottenhoff-Kalff <i>et al.</i> , Characterization of protein tyrosine kinases from human breast cancer: involvement of the c-src oncogene product, <i>Cancer Res</i> , 52(17):4773-4778 (1992)
116	Ozturk <i>et al.</i> , Detection of c-erbB-2 mRNAs using dig-labelled oligonucleotide probe with in situ hybridisation in human breast carcinoma: comparison with immunohistochemical results, <i>Anal Cell Pathol</i> , 16(4):201-209 (1998)
117	Porter <i>et al.</i> , Role of estrogen receptor/Sp1 complexes in estrogen-induced heat shock protein 27 gene expression, <i>Mol Endocrinol</i> , 10(11):1371-1378 (1996)
118	Qian and Lubman, Analysis of tryptic digests using microbore HPLC with an ion trap storage/reflectron time-of-flight detector, <i>Anal Chem</i> , 67(17):2870-2877 (1995)
119	Rodriguez <i>et al.</i> , Towards stoichiometric silver staining of proteins resolved in complex two-dimensional electrophoresis gels: real-time analysis of pattern development, <i>Electrophoresis</i> , 14(7):628-637 (1993)
120	Rosen <i>et al.</i> , Analysis of pp60c-src protein kinase activity in human tumor cell lines and tissues, <i>J Biol Chem</i> , 261(29):13754-13759 (1986)

Examiner:

Date Considered:

EXAMINER: Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

FORM PTO-1449
(Modified)U.S. Department of Commerce
Patent and Trademark Office

Attorney Docket No.: UM-06102

Serial No.: 09/778,548

INFORMATION DISCLOSURE STATEMENT BY APPLICANT
(Use Several Sheets if Necessary)Applicant: David M. Lubman *et al.*

Filing Date: 02/07/01

Group Art Unit:

(37 CFR § 1.98(b))

OTHER DOCUMENTS (Including Author, Title, Date, Relevant Pages, Place of Publication)

- | | |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 121 | Ruddon, Cancer biology, Oxford University Press, 3rd, 318-340 (1995) |
| 122 | Russell <i>et al.</i> , Cyclin D1 and D3 associate with the SCF complex and are coordinately elevated in breast cancer, <i>Oncogene</i> , 18(11):1983-1991 (1999) |
| 123 | Santner <i>et al.</i> , MCF-10CA1 cell lines: New highly tumorigenic derivatives of the MCF-10AT system, <i>Proc Am Assoc Cancer Res</i> , 39(202) (1998) (abstract only) |
| 124 | Schweinfest and Pappas, Subtraction hybridization: an approach to the isolation of genes differentially expressed in cancer and other biological system (Review), <i>Intern J Oncol</i> , 1(499-506 (1992) |
| 125 | Shekhar <i>et al.</i> , Transcriptional activation of functional endogenous estrogen receptor gene expression in MCF10AT cells: a model for early breast cancer, <i>Int J Oncol</i> , 13(5):907-915 (1998) |
| 126 | Shevchenko <i>et al.</i> , Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels, <i>Anal Chem</i> , 68(5):850-858 (1996) |
| 127 | Soule <i>et al.</i> , Isolation and characterization of a spontaneously immortalized human breast epithelial cell line, MCF-10, <i>Cancer Res</i> , 50(18):6075-6086 (1990) |
| 128 | Sturtevant, Applications of differential-display reverse transcription-PCR to molecular pathogenesis and medical mycology, <i>Clin Microbiol Rev</i> , 13(3):408-427 (2000) |
| 129 | Tesarik <i>et al.</i> , Estradiol modulates breast cancer cell apoptosis: a novel nongenomic steroid action relevant to carcinogenesis, <i>Steroids</i> , 64(1-2):22-27 (1999) |
| 130 | Tetu <i>et al.</i> , Prognostic significance of heat-shock protein-27 in node-positive breast carcinoma: an immunohistochemical study, <i>Breast Cancer Res Treat</i> , 36(1):93-97 (1995) |
| 131 | Thomas <i>et al.</i> , Exogenous estrogens and other risk factors for breast cancer in women with benign breast diseases, <i>J Natl Cancer Inst</i> , 69(5):1017-1025 (1982) |
| 132 | Williams <i>et al.</i> , Analysis of differential protein expression in normal and neoplastic human breast epithelial cell lines, <i>Electrophoresis</i> , 19(2):333-343 (1998) |
| 133 | Yang and Lee, Capillary isoelectric focusing-electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry for protein characterization, <i>Anal Chem</i> , 70(15):3235-3241 (1998) |
| 134 | Figeys and Aebersold, Nanoflow solvent gradient delivery from a microfabricated device for protein identifications by electrospray ionization mass spectrometry, <i>Anal Chem</i> , 70(18):3721-3727 (1998) (abstract only); |
| 135 | Figeys <i>et al.</i> , Data-dependent modulation of solid-phase extraction capillary electrophoresis for the analysis of complex peptide and phosphopeptide mixtures by tandem mass spectrometry: application to endothelial nitric oxide synthase, <i>Anal Chem</i> , 71(13):2279-2287 (1999) (abstract only); |
| 136 | Figeys <i>et al.</i> , Protein identification by solid phase microextraction-capillary zone electrophoresis-microelectrospray-tandem mass spectrometry, <i>Nat Biotechnol</i> , 14(11):1579-1583 (1996) (abstract only); |
| 137 | Figeys <i>et al.</i> , "An integrated microfluidics-tandem mass spectrometry system for automated protein analysis," <i>Anal. Chem.</i> 70:3728 (1998) (Abstract only); |
| 138 | Figeys <i>et al.</i> , Microfabricated device coupled with an electrospray ionization quadrupole time-of-flight mass spectrometer: protein identifications based on enhanced-resolution mass spectrometry and tandem mass spectrometry data, <i>Rapid Commun Mass Spectrom</i> , 12(20):1435-1444 (1998) (Abstract only); |
| 139 | Figeys <i>et al.</i> , A microfabricated device for rapid protein identification by microelectrospray ion trap mass spectrometry, <i>Anal Chem</i> , 69(16):3153-3160 (1997); |
| 140 | Figeys <i>et al.</i> , Optimization of solid phase microextraction - capillary zone electrophoresis - mass spectrometry for high sensitivity protein identification, <i>Electrophoresis</i> , 19(13):2338-2347 (1998) (abstract only) |
| 141 | Hunt <i>et al.</i> , Characterization of peptides bound to the class I MHC molecule HLA-A2.1 by mass spectrometry, <i>Science</i> . 1992 Mar 6;255(5049):1261-3 (abstract only) |
| 142 | Flyer, <i>et al.</i> , Identification by Mass Spectrometry of CD8+ T-Cell Mycobacterium tuberculosis Epitopes within the Rv0341 Gene Product, <i>Infection and Immunity</i> , 70(6):2926-2932, (June 2002); |
| 143 | Henderson <i>et al.</i> , HLA-A2.1-associated peptides from a mutant cell line: a second pathway of antigen presentation, <i>Science</i> . 1992 Mar 6;255(5049):1264-6 (abstract only) |

Examiner:

Date Considered:

EXAMINER:

Initial citation considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.